

On Tubular vs. Swung surfaces

Tomás Recio^a, J. Rafael Sendra^{b,*}, Luis-Felipe Tabera^a, Carlos Villarino^b

^a*Departamento de Matemáticas Universidad de Cantabria, 39071, Santander, Spain*

^b*Dpto. de Física y Matemáticas, Universidad de Alcalá E-28871, Alcalá de Henares, Madrid, Spain*

Abstract

We determine necessary and sufficient conditions for a tubular surface to be swung, and viceversa. From these characterizations, we derive two symbolic algorithms. The first one decides whether a given implicit equation, of a tubular surface, admits a swung parametrization and, in the affirmative case, it outputs such a parametrization. The second one decides whether a given swung surface parametrization is a tubular surface and, in the affirmative case, it outputs the implicit equation.

Keywords: Swung surface, tubular surface.

1. Introduction

2 In CAGD, many different families of surfaces are usually considered. For
3 instance, we may talk about revolution, ruled, tubular, swung, swept, etc. sur-
4 faces (see [7] for a nice survey on these different families of surfaces). However,
5 it is possible that a surface belongs to more than one of these families. For
6 example, every revolution surface is an instance of a swung surface, that is also
7 an example of swept surface.

8 But the inclusion of different families of surfaces into each other, does not
9 hold in general. In many cases, surfaces belonging to a particular family have
10 to verify some extra conditions in order to be, as well, members of a different
11 family of surfaces. These extra conditions usually take an algebraic form, so
12 the intersection of the two families of surfaces has measure zero in the space
13 representing each family. But, when this happens, the manipulation of such a
14 surface, belonging to several families of surfaces, can profit from the accumu-
15 lated knowledge about surfaces on each concrete family. For instance, elements
16 belonging to some families can have a simple implicit description, while those
17 pertaining to some other families could enjoy having straightforward paramet-
18 ric representations. Belonging simultaneously to two families of such different

*Corresponding author. Tel: +34 918854902. Fax: +34 918854951.

Email addresses: tomas.recio@unican.es (Tomás Recio), rafael.sendra@uah.es (J. Rafael Sendra), luisfelipe.tabera@unican.es (Luis-Felipe Tabera), carlos.villarino@uah.es (Carlos Villarino)

19 kinds could result, for example, in an easier method for solving the symbolic
20 implicit/parametric conversion for the given surface.

21 In this paper, we determine those surfaces that are simultaneously tubular
22 and swung.

23 Tubular surfaces are those irreducible surfaces described by an implicit equa-
24 tion

$$A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$$

25
26 where $A, B, C \in \mathbb{R}[x_3]$, $\gcd(A, B, C) = 1$ and the total degree w.r.t. $\{x_1, x_2\}$
27 is 2; note that in a tubular surface it cannot happen that two of the poly-
28 nomials A, B, C vanish simultaneously. Notice that any surface with a pencil
29 of rational curves is birational equivalent to a tubular surface. Algorithms to
30 parametrize a tubular surface are described in [4], where it is also shown that
31 many instances of the real surface parametrization problem can be reduced to
32 the tubular case. See [2], Example 2.3 for an application in the context of swung
33 surfaces. The importance of tubular surfaces concerning this relevant, generally
34 unsolved, problem of parametrizing over the reals, is one of the reasons for our
35 choice of tubular surfaces as one of the families in our double test approach.

36 On the other hand, swung surfaces are a generalization of the well known
37 revolution surfaces (around the x_3 -axis) in which a profile curve parametrized by
38 $(0, \phi_1(t), \phi_2(t))$ is transported *around* a trajectory curve $(\psi_1(s), \psi_2(s), 0)$. The
39 obtained surface is the surface parametrized by

$$(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t)).$$

40
41 If the trajectory curve is a circle, then the swung surface is just the classical
42 revolution surface. Swung surfaces have been subject of recent research, even
43 considering elementary issues as the problem of implicitizing; see [6] where the
44 authors use μ -bases to develop specific techniques for implicitation of swung
45 surfaces, as an alternative of the well-know techniques in elimination theory.

46 Notice that, if the profile curve of a revolution surface is given by the graph
47 of a rational function $x_2 = (f/g)(x_3)$, then the revolution surface has equation
48 $g(x_3)^2x_1^2 + g(x_3)^2x_2^2 - f(x_3)^2 = 0$ and this is clearly a tubular surface. Conversely,
49 a necessary condition for a swung surface to be tubular is that its intersection
50 with the family of planes $\{x_3 = c\}$ is a pencil of conics. However, this is not
51 sufficient in general, see Example 4.2.

52 Since tubular surfaces are very relevant in the real reparametrization prob-
53 lem and swung surfaces are very useful in CAD, in this paper we want to merge
54 both advantages and determine necessary and sufficient conditions for a tubular
55 surface to be swung and viceversa (Theorems 2.3 and 3.1). These characteriza-
56 tions provide a symbolic algorithm that passes from an implicit tubular repre-
57 sentation to a swung parametrization whenever possible, as well as a symbolic
58 algorithm that decides whether a swung parametrization is tubular and, if so, it
59 computes the implicit equation. An example of the above mentioned advantages
60 of being simultaneously in both categories is developed in Example 4.3.

61 Throughout the paper we assume that the implicit representations of the

62 tubular surfaces are real polynomials, and that the swung surface parametriza-
63 tions are real.

64 2. From tubular to swung

65 We show how to decide if a tubular surface is swung and, then, how to
66 compute a swung parametrization. A naive approach could start applying
67 parametrization algorithms to the given implicit equation of the tubular sur-
68 face, expecting to obtain swung parametrization (if the surface is swung). But
69 parametrization algorithms are not trivial and, even if a parametrization is ob-
70 tained, it is not expected that it will have the structure of a swung parametriza-
71 tion. Thus, we need to develop some specific techniques to deal with this prob-
72 lem.

73 We start this section analyzing some special cases. We already know that
74 two of the polynomials A, B, C can not vanish simultaneously. Let us study
75 what happens when one of them vanishes.

76 **Lemma 2.1.** *Let $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$ be the implicit equation of a*
77 *tubular surface. If $AB = 0$, and the surface is rational over \mathbb{R} , then it is swung.*

78 *Proof.* By definition of tubular surface, we know that A, B cannot be simul-
79 taneously zero. Let $A = 0$ but $B \neq 0$. Let $P(u, v) = (u, M(v), N(v))$ be a
80 proper real parametrization of the tubular surface $B(x_3)x_2^2 + C(x_3) = 0$. We
81 observe that M is not zero, because the surface is not a plane. Then, taking
82 $Q(s, t) = P(M(t)s, t)$ we get $Q(s, t) = (M(t)s, M(t), N(t))$ that is a swung
83 proper parametrization (note that $(M(t)s, t)$ is a birational map of \mathbb{R}^2 on \mathbb{R}^2)
84 with profile curve $(0, M(t), N(t))$ and trajectory curve $(s, 1, 0)$.

85 Let $A \neq 0$ but $B = 0$. Then, the same reasoning works. In this case
86 if $P(u, v) = (M(v), u, N(v))$ then the profile curve is $(0, M(t), N(t))$ and the
87 trajectory curve is $(1, s, 0)$. \square

88 **Lemma 2.2.** *Let $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$ be the implicit equation of a*
89 *tubular surface. If $C = 0$ the surface is not swung.*

90 *Proof.* Assume that $(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t))$ is a swung parametrization
91 of the surface. Then

$$92 \quad \phi_1(t)^2(A(\phi_2(t))\psi_1(s)^2 + B(\phi_2(t))\psi_2(s)^2) = 0.$$

93 Since $\phi_1(t)$ is not zero, because otherwise the surface would degenerate to a
94 curve, then

$$95 \quad A(\phi_2(t))\psi_1(s)^2 + B(\phi_2(t))\psi_2(s)^2 = 0.$$

96 In addition, $AB \neq 0$. So, $A(\phi_2(t))B(\phi_2(t)) \neq 0$. On the other hand, we also
97 have that not both rational functions $\psi_i(s)$ can be zero; say $\psi_1(s) \neq 0$. Then

$$98 \quad \frac{A(\phi_2(t))}{B(\phi_2(t))} = -\frac{\psi_2(s)^2}{\psi_1(s)^2}.$$

99 This implies that $A(\phi_2(t)) = \lambda B(\phi_2(t))$ for some $\lambda \in \mathbb{R} \setminus \{0\}$. Thus, $A(x_3) =$
100 $\lambda B(x_3)$. Moreover, since $\gcd(A, B, C) = 1$ then A and B must be constants, and
101 the equation of the tubular surface is $\lambda x_1^2 + x_2^2 = (x_2 - \sqrt{-\lambda}x_1)(x_2 + \sqrt{-\lambda}x_1) = 0$.
102 But this is a contradiction, because a tubular surface is irreducible. \square

103 Taking into account the previous lemmas we will assume that $ABC \neq 0$.

104 **Theorem 2.3.** *Let $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$ be the implicit equation of*
105 *a real tubular surface, such that $ABC \neq 0$, $\gcd(A, B, C) = 1$. Then, the surface*
106 *is a swung surface if and only if:*

- 107 1. $B(x_3)/A(x_3) = k \in \mathbb{R}$ is constant.
- 108 2. One of the curves (or a component of) $A(y)x^2 \pm C(y)$ is rational parametriz-
109 able over \mathbb{R} .

110 *Proof.* Assume that $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$ is swung. Then, there exists
111 a swung parametrization $(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t))$ of the surface. Hence

$$112 \quad A(\phi_2(t))\phi_1(t)^2\psi_1(s)^2 + B(\phi_2(t))\phi_1(t)^2\psi_2(s)^2 + C(\phi_2(t)) = 0.$$

113 We observe that $\phi_2(t)$ cannot be a constant, because the surface is not a plane.
114 Also, note that $\phi_1(t)$ cannot be zero, since otherwise the given variety would
115 be a line. This, in particular implies that $C(\phi_2(t))B(\phi_2(t))A(\phi_2(t))\phi_1(t)$ is not
116 zero. So, manipulating the above expression, we get that

$$117 \quad \frac{\psi_1(s)^2}{\alpha(t)} + \frac{\psi_2(s)^2}{\beta(t)} - 1 = 0,$$

118 where

$$119 \quad \alpha(t) = -\frac{C(\phi_2(t))}{A(\phi_2(t))\phi_1(t)^2} \neq 0, \quad \beta(t) = -\frac{C(\phi_2(t))}{B(\phi_2(t))\phi_1(t)^2} \neq 0. \quad (1)$$

120 Therefore $(\psi_1(s), \psi_2(s))$ parametrizes the conic, defined over $\mathbb{R}(t)$ by $x_1^2/\alpha(t) +$
121 $x_2^2/\beta(t) = 1$. However, since $(\psi_1(s), \psi_2(s))$ is over \mathbb{R} , its implicit equation is
122 over \mathbb{R} . So, since $x_1^2/\alpha(t) + x_2^2/\beta(t) = 1$ is irreducible as conic over $\mathbb{R}(t)$, we
123 get that both implicit equations must be equal, and hence $\alpha(t), \beta(t) \in \mathbb{R} \setminus \{0\}$.
124 Thus, $\alpha(t)/\beta(t) = (B/A)(\phi_2(t))$ is constant. Hence, since ϕ_2 is not constant,
125 $B(x_3)/A(x_3) = k \in \mathbb{R} \setminus \{0\}$ is constant.

126 Moreover, by equation (1), $(\phi_1(t), \phi_2(t))$ is a parametrization (of a compo-
127 nent of) the curve defined by $C(y) + \alpha x^2 A(y)$ (recall that $\alpha(t) \in \mathbb{R} \setminus \{0\}$) and
128 $(\phi_1/\sqrt{|\alpha|}, \phi_2)$ is a parametrization (of a component of) $C(y) + \text{sign}(\alpha)x^2 A(y)$.

129 Assume now that we have a tubular surface

$$130 \quad A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3)$$

131 such that $ABC \neq 0$, $B/A = k \in \mathbb{R}$ is constant and that $(\phi_1(t), \phi_2(t))$ is a real
132 parametrization of (a component of) $C(y) \pm x^2 A(y)$. We want to prove that it
133 is swung. Consider the profile curve $(0, \phi_1(t), \phi_2(t))$. We have to construct a
134 sliding curve adapted to the tubular surface.

135 Consider the conic $x_1^2 + kx_2^2 = \pm 1$. This will be our trajectory curve; note
 136 that $k \neq 0$. Let $(\psi_1(s), \psi_2(s))$ be a parametrization of the conic (we will see
 137 below that the parametrization can be taken over \mathbb{R}), we have to prove that
 138 $(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t))$ parametrizes the surface. But

$$139 \quad A(\phi_2(t))\phi_1(t)^2\psi_1(s)^2 + B(\phi_2(t))\phi_1(t)^2\psi_2(s)^2 + C(\phi_2(t)) =$$

$$140 \quad A(\phi_2(t))\phi_1(t)^2(\psi_1(s)^2 + k\psi_2(s)^2) + C(\phi_2(t)) = \pm A(\phi_2(t))\phi_1(t)^2 + C(\phi_2(t)) = 0.$$

142 It only remains to prove that the corresponding conic $x_1^2 + kx_2^2 = \pm 1$ is real, from
 143 where it follows that the parametrization $(\psi_1(s), \psi_2(s))$ can always be taken over
 144 \mathbb{R} . For this purpose, we distinguish several cases. We first observe that, since
 145 $\gcd(A, B, C) = \gcd(A, C) = 1$, if $C(y) \pm x^2A(y)$ factors then it has two factors
 146 and they are linear in x , and hence both rational. Let \mathcal{C}^\pm be the curve defined
 147 by $C(y) \pm x^2A(y)$. Furthermore, we note that \mathcal{C}^+ (resp. a component of it) is
 148 rational (over \mathbb{C}) iff \mathcal{C}^- (resp. a component of it) is rational (over \mathbb{C}).

149 (i) Let \mathcal{C}^+ (or a component of it) be parametrizable over \mathbb{R} . Then we have
 150 to parametrize $x_1^2 + kx_2^2 = 1$ that is always real, independently of the sign
 151 of k .

152 (ii) Let \mathcal{C}^+ (nor a component of it) not be parametrizable over \mathbb{R} . Then, by
 153 hypothesis, \mathcal{C}^- (or a component of it) is parametrizable over \mathbb{R} . In this
 154 case, we have to parametrize $x_1^2 + kx_2^2 = -1$. We prove that $k < 0$ and, so,
 155 the conic real. Let us assume that $k > 0$. No component of \mathcal{C}^+ is a real
 156 curve. Therefore, the curve \mathcal{C}^+ cannot have a real regular point. On the
 157 other hand, the tubular surface, that is defined by $A(y)(x_1^2 + kx_2^2) + C(y)$,
 158 is a real surface. Therefore, it contains a regular real point $P = (\alpha, \beta, \gamma)$.
 159 So,

$$160 \quad A(\gamma)(\alpha^2 + k\beta^2) + C(\gamma) = 0 \quad (2)$$

Observe that $A(\gamma) \neq 0$, since otherwise $C(\gamma) = 0$ and $\gcd(A, B, C) \neq 1$
 which is a contradiction. Now, since P is regular, we have that either
 $\alpha A(\gamma) \neq 0$ or $k\beta A(\gamma) \neq 0$ or $A'(\gamma)(\alpha^2 + k\beta^2) + C'(\gamma) \neq 0$. That is (note
 that $k \neq 0$) either $\alpha \neq 0$ or $\beta \neq 0$ or $A'(\gamma)(\alpha^2 + k\beta^2) + C'(\gamma) \neq 0$. In
 addition, since $A(\gamma) \neq 0$ we have that

$$Q := \left(\pm \sqrt{-\frac{C(\gamma)}{A(\gamma)}}, \gamma \right) \in \mathcal{C}^+.$$

161 We analyze each case.

162 – Let $\alpha \neq 0$. We observe that $C(\gamma) \neq 0$ because: if $C(\gamma) = 0$, since
 163 $A(\gamma) \neq 0$, by (2), one has that $\alpha^2 + k\beta^2 = 0$ but this is impossible
 164 because $\alpha \neq 0$ and $k > 0$. But this implies that the square of the
 165 partial derivative w.r.t. x of $C(y) + x^2A(y)$ at Q is $-4C(\gamma)A(\gamma) \neq 0$.
 166 Thus Q is a regular point of \mathcal{C}^+ . Therefore, Q cannot be real. So
 167 $C(\gamma)A(\gamma) > 0$. But, from (2), we get then that $\alpha^2 + k\beta^2 < 0$ which
 168 is impossible since $\alpha \neq 0$ and $k > 0$.

- 169 – If $\beta \neq 0$ the reasoning is above.
 170 – Let $A'(\gamma)(\alpha^2 + k\beta^2) + C'(\gamma) \neq 0$. Because of the two previous cases,
 171 we can assume w.l.o.g. that $\alpha = \beta = 0$. So, by (2), we have that
 172 $C'(\gamma) = 0$. So $Q = (0, \gamma)$. Then, the derivative w.r.t. y of $C(y) +$
 173 $x^2A(y)$ at Q is $C'(\gamma)$. Therefore, since Q is real, we have that $C'(\gamma) =$
 174 0 which contradicts our assumption.

175 □

176 **Remark 2.4.** Assume $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0$ is a tubular surface with
 177 a swung parametrization. Then by the previous result, $B(x_3)/A(x_3) = k \in \mathbb{R}$.
 178 So, since C is not zero (see Lemma 2.2), all non-degenerated sections with the
 179 family of planes $\{x_3 = c\}$ will yield conics of the same type, either they are
 180 all ellipses or all hyperbolas. When $A = 0$ or $B = 0$, sections with $x_3 = c$
 181 degenerate to a pair of lines.

182 Lemma 2.1 and Theorem 2.3 show how to check whether a rational tubular
 183 surface is swung. Moreover, their proofs provide a method to find a swung
 184 parametrization of a swung tubular surface. More precisely, one has the follow-
 185 ing algorithm.

186 (Parametrization) Algorithm Tubular/Swung

187 Input: let $A(x_3)x_1^2 + B(x_3)x_2^2 + C(x_3) = 0, C \neq 0, \gcd(A, B, C) = 1$, be the
 188 implicit equation of a rational tubular surface \mathcal{S} .

189 Output: decision on whether \mathcal{S} admits a swung parametrization or not. If so, a
 190 swung parametrization of \mathcal{S} is obtained.

- 191 1. If $A = 0$, compute a real parametrization (M, N) of $B(x_3)x_2^2 + C(x_3) = 0$
 192 (see [3]), and return $(M(t)s, M(t), N(t))$ as parametrization of the surface.
- 193 2. If $B = 0$, compute a real parametrization (M, N) of $A(x_3)x_1^2 + C(x_3) =$
 194 0 (see [3]), and return $(M(t), M(t)s, N(t))$ as a parametrization of the
 195 surface.
- 196 3. Else (i.e. $AB \neq 0$)
 - 197 (a) Compute $k = B/A$. If k is not constant then return that \mathcal{S} is not
 198 swung.
 - 199 (b) [Profile curve computation] Apply algorithm in [3] to compute a real
 200 parametrization $(\psi_1(t), \psi_2(t))$ of (a factor of) one of the curves $C \pm$
 201 x^2A ; if no component is parametrizable over \mathbb{R} then return that \mathcal{S}
 202 is not swung. Take $\epsilon = 0$ if $(\phi_1(t), \phi_2(t))$ parametrizes (a factor of)
 203 $C + x^2A$ and $\epsilon = 1$ if parametrizes (a factor of) $C - x^2A$.
 - 204 (c) [Trajectory curve computation] Compute a real parametrization $(\psi_1(s),$
 205 $\psi_2(s))$ of the conic $x^2 + ky^2 = (-1)^\epsilon$.

206 **3. From swung to tubular**

207 We now work the other way around, given a swung surface, detect if it
 208 is tubular. Of course, one could just implicitize the surface and check if the

209 implicit equation corresponds to a tubular surface, but we will provide a char-
 210 acterization based on the profile and trajectory curves, providing insight and
 211 not simply blind, costly, resultant or Gröbner bases implicitization algorithms.
 212 See Example 4.1.

213 **Theorem 3.1.** *Let $(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t))$ be a parametrization of a*
 214 *swung surface different from the planes $x_1 = 0, x_2 = 0$. The surface is tubular*
 215 *if and only if*

- 216 1. *The trajectory curve $(\psi_1(s), \psi_2(s))$ is either a conic in normal form $x_1^2/a +$*
 217 *$x_2^2/b - 1 = 0$ or a line of the form $x_1 = \lambda$ or $x_2 = \lambda$, with $\lambda \neq 0$.*
- 218 2. *There exists a rational function h such that $\phi_1(t)^2 = h(\phi_2(t))$.*

219 *Proof.* Assume that the surface is tubular. There exists A, B, C such that

$$220 \quad A(\phi_2)\phi_1^2\psi_1^2 + B(\phi_2)\phi_1^2\psi_2^2 + C(\phi_2) = 0$$

221 We distinguish cases. Let $AB \neq 0$. By Theorem 2.3, the surface is tubu-
 222 lar and swung, so $B(x_3) = kA(x_3)$. Evaluating at a value $t = t_0$, we get
 223 that $(\psi_1(s), \psi_2(s))$ parametrizes the curve $x_1^2/a + kx_2^2/a - 1 = 0$, where $a =$
 224 $-C(\phi_2(t_0))/(A(\phi_2(t_0))\phi_1(t_0)^2)$. Now, manipulating the equation we get

$$\begin{aligned} 225 \quad 0 &= A(\phi_2)\phi_1^2\psi_1^2 + B(\phi_2)\phi_1^2\psi_2^2 + C(\phi_2) = \\ 226 \quad &= A(\phi_2)\phi_1^2(\psi_1^2 + k\psi_2^2) + C(\phi_2) = \\ 227 \quad &= A(\phi_2)\phi_1^2a + C(\phi_2) \end{aligned}$$

230 So, $\phi_1^2 = -C(\phi_2)/(aA(\phi_2)) = h(\phi_2)$.

231 Let $A = 0$, then $BC \neq 0$. Evaluating at a value $t = t_0$, we get that
 232 $B(\phi_2(t_0))\phi_1(t_0)^2\psi_2(s)^2 + C(\phi_2(t_0)) = 0$. So $\psi_2(s)$ is constant, say λ , and the
 233 trajectory curve is the line $x_2 = \lambda$; note that, since ψ_2 is real then $\lambda \in \mathbb{R}$ and
 234 since the surface is not $x_2 = 0$ then $\lambda \neq 0$. Moreover, $\phi_1^2 = -C(\phi_2)/(\lambda^2B(\phi_2)) =$
 235 $h(\phi_2)$. If $B = 0$ the reasoning is similar.

236 Conversely, let us assume first that (ψ_1, ψ_2) parametrizes a conic $x_1^2/a +$
 237 $x_2^2/b - 1$ and that $\phi_1^2 = C(\phi_2)/A(\phi_2)$ for some $a, b \in \mathbb{R}, C, A \in \mathbb{R}[x_3]$, with
 238 $\gcd(A, C) = 1$ and $C \neq 0$ (note that ϕ_1 cannot be zero). We want to prove that
 239 the given surface is tubular. Consider the equation

$$240 \quad A(x_3)x_1^2 + a/bA(x_3)x_2^2 - aC(x_3) = 0$$

241 Substituting the parametrization, we get

$$242 \quad A(\phi_2)\phi_1^2(\psi_1^2 + a/b\psi_2^2) + aC(\phi_2) = aA(\phi_2)\phi_1^2 - aC(\phi_2) = 0$$

243 so the surface is tubular. Note that, by construction $\gcd(A, a/bA, C) = 1$ and
 244 the total degree w.r.t. $\{x_1, x_2\}$ is 2 because C is no zero. Now, let us assume
 245 that (ψ_1, ψ_2) parametrizes a line $x_2 = \lambda$, and that $\phi_1^2 = C(\phi_2)/B(\phi_2)$ for some

246 $\lambda \in \mathbb{R} \setminus \{0\}$, $C, B \in \mathbb{R}[x_3]$, with $\gcd(B, C) = 1$ and $C \neq 0$. Then, the surface is
 247 the tubular surface of equation

$$248 \quad \frac{1}{\lambda^2} B(x_3)x_2^2 - C(x_3) = 0.$$

249 If the trajectory curve is a line of the type $x_1 = \lambda$, with $\lambda \neq 0$, the reasoning is
 250 similar.

251 □

252 **Remark 3.2.** In order to compute polynomials A and C such that $\phi_1^2 =$
 253 $C(\phi_2)/A(\phi_2)$, we may use rational decomposition techniques [1]. In particu-
 254 lar if $n = \max\{\deg(\text{numer}(\phi_2)), \deg(\text{denom}(\phi_2))\}$, $m = \max\{\deg(\text{numer}(\phi_1)),$
 255 $\deg(\text{denom}(\phi_1))\}$, then the degree of A and C is bounded by $2m/n$. If $2m/n$ is
 256 not an integer then there is no solution and the surface is not tubular. If $2m/n$
 257 is an integer, we may take A and C as polynomials of degree $2m/n$ with un-
 258 determined coefficients, then evaluate the expression $\phi_1(t)^2 C(\phi_2(t)) = A(\phi_2(t))$
 259 at $4m/n + 2$ values of t where $C(\phi_2(t)) \neq 0$ and, finally compute the coefficients
 260 of A and C by solving the resulting linear system of equations.

261 Using the last argument in the previous remark, we get the following corol-
 262 laries of Theorem 3.1.

263 **Corollary 3.3.** *If the (implicit) profile curve of a swung surface has degree*
 264 *bigger than 2 w.r.t. the first variable, then it is not tubular.*

265 Using Theorem 4.2.1. in [5], one has the next result

266 **Corollary 3.4.** *Let $(0, \phi_1, \phi_2)$, with $\phi_2 \neq 0$, be a proper parametrization of the*
 267 *profile curve of a swung surface. If $\deg(\phi_2) > 2$ then the surface is not tubular.*

268 We finish the section with an algorithm that decides whether a swung surface
 269 is tubular and, in the affirmative case, computes the implicit equation.

270 **(Implicitization) Algorithm Swung/Tubular**

271 **Input:** Let $(\phi_1(t)\psi_1(s), \phi_1(t)\psi_2(s), \phi_2(t))$ be a parametrization of a swung surface
 272 \mathcal{S} different from the planes $x_1 = 0, x_2 = 0$.

273 **Output:** decision on whether \mathcal{S} is tubular or not. If \mathcal{S} is tubular the implicit
 274 equation is also obtained

275 1. Check whether $(\psi_1(s), \psi_2(s))$ is in one of the following cases

276 (i) it is a conic in normal form $x_1^2/a + x_2^2/b - 1 = 0$.

277 (ii) it is a line of the form $x_1 = \lambda$, with $\lambda \neq 0$.

278 (iii) it is a line of the form $x_2 = \lambda$, with $\lambda \neq 0$.

279 If the answer is no, then return that \mathcal{S} is not tubular.

280 2. Use Remark 3.2 to check whether there exists a rational function $h = C/A$
 281 with $\gcd(A, C) = 1, C \neq 0$, such that $\phi_1(t)^2 = h(\phi_2(t))$. If the answer is
 282 yes, then return

- 283 (a) If in Step 1, (i) holds then return that \mathcal{S} is tubular and that $A(x_3)x_1^2 +$
 284 $\frac{a}{b}A(x_3)x_2^2 - aC(x_3)$ is its implicit equation.
 285 (b) If in Step 1, (ii) holds then return that \mathcal{S} is tubular and that $\frac{1}{\lambda^2}A(x_3)x_1^2 -$
 286 $C(x_3)$ is its implicit equation.
 287 (c) If in Step 1, (iii) holds then return that \mathcal{S} is tubular and that $\frac{1}{\lambda^2}A(x_3)x_2^2 -$
 288 $C(x_3)$ is its implicit equation.
 289 If the answer is no, then return that \mathcal{S} is not tubular else return that \mathcal{S} is
 290 tubular.

291 4. Examples

292 We illustrate our results by some examples.

293 **Example 4.1.** We consider the swung surface (see Fig. 1)

$$294 \left(4 \frac{(t^2 + t + 1)(s^2 - 1)}{(t^3 + 2)(s^2 + 1)}, 18 \frac{(t^2 + t + 1)s}{(t^3 + 2)(s^2 + 1)}, t \right).$$

Without any computation, it is elementary to see that the trajectory curve




Figure 1: Tubular, swung, surface in Example 4.1

295 is the ellipse $(1/16)x^2 + (1/81)y^2 - 1$ (step 1.i from the above algorithm); so
 296 $a = 16, b = 81$, and $h = (z^2 + z + 1)^2 / (z^3 + 2)^2$ (step 2). Therefore the surface
 297 is tubular and its implicit equation is (step 2.a)
 298

$$299 (x_3^3 + 2)^2 x_1^2 + \frac{16}{81} (x_3^3 + 2)^2 x_2^2 - 16 (x_3^2 + x_3 + 1)^2$$

300 **Example 4.2.** Consider the swung surface defined by (see Fig. 2)

$$301 \left(t^2 \frac{s^2 - 1}{s^2 + 1}, t^2 \frac{2s}{s^2 + 1}, t^3 \right).$$




Figure 2: Non-Tubular, swung, surface in Example 4.2

Since t^4 cannot be expressed as $h(t^3)$, the surface is not tubular (step 2). Observe also that the profile curve is $(0, t^2, t^3)$ that is proper and the degree of ϕ_2 is $3 > 2$ (see Corollary 3.4). However, for any value of $t = t_0$, excluding 0, the curve

$$\left(t_0^2 \frac{s^2 - 1}{s^2 + 1}, t_0^2 \frac{2s}{s^2 + 1}, t_0^3 \right)$$

302 is a circle. The implicit equation of the surface is $x_1^6 + 3x_1^4x_2^2 + 3x_1^2x_2^4 + x_2^6 - x_3^4 = 0$.
 303 We may express this polynomial as

$$304 \quad \left(x_1^2 + x_2^2 - \sqrt[3]{x_3^4} \right) \left(x_1^2 + x_2^2 + \frac{1 - i\sqrt{3}}{2} \sqrt[3]{x_3^4} \right) \left(x_1^2 + x_2^2 + \frac{1 + i\sqrt{3}}{2} \sqrt[3]{x_3^4} \right)$$

305 and notice that the pencil of conics $\left(x_1^2 + x_2^2 - \sqrt[3]{x_3^4} \right)$ is not rational.

306 **Example 4.3.** $F \equiv (-36)x^2 + (-32z)xy + (4z^2 - 100)y^2 + (16z^2 + 144)x +$
 307 $(-4z^3 + 164z)y + z^4 - 82z^2 + 81 = 0$. This surface (See Fig. 3) is a pencil of
 308 conics and, can be transformed into a tubular surface. In fact, let us take the
 309 following $\mathbb{R}(z)$ -change of variables $x_1 = x + (\frac{4}{9}z)y - \frac{2}{9}z^2 - 2$, $y_1 = y - 1/2z$. Then
 310 F is transformed into $F^* \equiv -36x_1^2 + (100/9z^2 - 100)y_1^2 + (-25z^2 + 225) = 0$
 311 (see Fig. 4 left). This surface is not in the hypotheses of Theorem 2.3, so it
 312 is tubular but not swung (step 3.a of the Tubular/Swung algorithm). Now,
 313 consider the new change of variables $x_2 = 1/x_1$, $y_2 = y_1/x_1$, we get the surface
 314 $F^{**} = (-25z^2 + 225)x_2^2 + (100/9z^2 - 100)y_2^2 - 36$ (Fig. 4 right), where $A =$
 315 $(-25z^2 + 225)$, $B = (100/9z^2 - 100)$, $C = -36$. This tubular surface verifies
 316 the hypotheses of Theorem 3.1, since $A/B = -9/4 = k$ (step 3.a) and the curve
 317 $A(y)x^2 + C = -25x^2y^2 + 225x^2 - 36$ is parametrizable by the profile curve (step




Figure 3: Surface F in Example 4.3

318 3.b):

$$319 \left(\frac{2t^2 + 1}{5t^2 - 1}, \frac{6t}{t^2 + 1} \right).$$

320 Following that theorem, the trajectory curve is $x_2^2 - 4/9y_2^2 - 1 = 0$ that we can
 321 parametrize as $\left(\frac{s^2+1}{s^2-1}, \frac{3s}{s^2-1}\right)$ (step 3.c). This provides the following parametriza-
 322 tion of the surface

$$323 \begin{cases} x_2 &= \frac{2}{5} \frac{t^2+1}{t^2-1} \cdot \frac{s^2+1}{s^2-1} \\ y_2 &= \frac{2}{5} \frac{t^2+1}{t^2-1} \cdot \frac{3s}{s^2-1} \\ z &= \frac{6t}{t^2+1} \end{cases}$$

324 Finally, reverting the change of variables, we get the following parametrization
 325 of the original surface

$$326 \begin{cases} X &= -\frac{4\phi_2\psi_2}{9\psi_1} + \frac{1}{\phi_1\psi_1} + 2 \\ Y &= \frac{1}{2}\phi_2 + \frac{\psi_2}{\psi_1} \\ Z &= \phi_2 \end{cases}$$

327 with $\phi_1 = \frac{2}{5} \frac{t^2+1}{t^2-1}$, $\phi_2 = \frac{6t}{t^2+1}$, $\psi_1 = \frac{s^2+1}{s^2-1}$, $\psi_2 = \frac{3s}{s^2-1}$.

$$328 \begin{cases} X &= \frac{1}{2} \cdot \frac{(3ts-t+s-3) \cdot (3ts+t-s-3)}{(s^2+1) \cdot (t^2+1)} \\ Y &= 3 \cdot \frac{(t+s) \cdot (ts+1)}{(s^2+1) \cdot (t^2+1)} \\ Z &= \frac{6t}{t^2+1} \end{cases}$$

329 **Acknowledgments:** Authors supported by the Spanish *Ministerio de Eco-*
 330 *nomía y Competitividad* and by the European Regional Development Fund




Figure 4: Left: Tubular, non-swung, surface F^* in Example 4.3. Right: Tubular, swung, surface F^{**} in Example 4.3

331 (ERDF), under the Project MTM2011-25816-C02-(01,02). Second and fourth
 332 authors are members of the research group ASYNACS (Ref. CCEE2011/R34)

333 Bibliography

- 334 [1] Cesar Alonso, Jaime Gutiérrez, and Tomás Recio. A rational function de-
 335 composition algorithm by near-separated polynomials. *J. Symbolic Comput.*,
 336 19(6):527–544, 1995.
- 337 [2] Carlos Andradas, Tomás Recio, J. Rafael Sendra, Luis-Felipe Tabera, and
 338 Carlos Villarino. Reparametrizing swung surfaces over the reals. *Appl.*
 339 *Algebra Engrg. Comm. Comput.*, 25(1-2):39–65, 2014.
- 340 [3] Tomás Recio and J. Rafael Sendra. Real reparametrizations of real curves.
 341 *J. Symbolic Comput.*, 23(2-3):241–254, 1997. Parametric algebraic curves
 342 and applications (Albuquerque, NM, 1995).
- 343 [4] Josef Schicho. Proper parametrization of real tubular surfaces. *J. Symbolic*
 344 *Comput.*, 30(5):583–593, 2000.
- 345 [5] J. Rafael Sendra, Franz Winkler, and Sonia Pérez-Díaz. *Rational algebraic*
 346 *curves: A computer algebra approach*, volume 22 of *Algorithms and Compu-*
 347 *tation in Mathematics*. Springer, Berlin, 2008.
- 348 [6] Xiaoran Shi and Ron Goldman. Implicitizing rational surfaces of revolution
 349 using μ -bases. *Comput. Aided Geom. Design*, 29(6):348–362, 2012.
- 350 [7] Yuan Zhao, Yan Zhou, John L. Lowther, and Ching-Kuang Shene. Cross-
 351 sectional design: a tool for computer graphics and computer-aided design
 352 courses. In *Conference on Frontiers in Education*, volume 2, 1999.