

Pyridinium-N-(2-pyridyl)aminides: A Selective Approach to Substituted 2-Aminopyridines


Rosa Carceller, Jose L. García-Navío, María L. Izquierdo and Julio Alvarez-Builla*

Departamento de Química Orgánica, Universidad de Alcalá de Henares, 28871 Alcalá de Henares, Madrid, España.

Abstract: Differently substituted 2-aminopyridines have been prepared in two steps from pyridinium-N-(2-pyridyl)aminide, by reaction with the corresponding electrophiles and reduction of the N-N bond.

As the body of knowledge about the reactivity of azinium *N*-ylides expands, the interest in such compounds as building blocks for the synthesis of heterocyclic derivatives continuously increases.¹ Some of our work in the field has been concerned with the synthesis of heteroaryl-stabilized cycloiminium ylides as a way of producing highly stable dipoles, with heterocyclic moieties of unusual reactivity.²

In this paper we wish to report our first results with the pyridinium-N-(2-pyridyl)aminide **1** in which there is a negatively charged 2-aminopyridine fragment. The precursor pyridinium bromide was prepared by using the method of Bayer,³ and converted into the stable aminide **1** by treatment with K_2CO_3 /acetone. The structure of **1** in solution showed an intramolecular hydrogen bond as indicated, which has been described for related compounds.⁴



Because of its structure in solution, reaction of **1** with alkyl halides produced regioselective alkylation on the exocyclic nitrogen (compounds **2**, Fig. 2), with no signs of alkylation on the N1 of the 2-pyridyl ring.

Reaction of **1** with electrophiles producing C-substitution was obtained under very mild conditions. Thus, iodination with equimolar amounts of iodine at room temp., yielded the monoiodo salt **3** (61%), while when iodination was made in the presence of base (K_2CO_3), the initially formed **3** was deprotonated and attacked again, being the diiodo betaine **4** isolated (22%). On treatment with bromine, the process was out of control, being obtained then a mixture of 5-bromo and 3,5-dibromo salts (1:4), even when working at $0^\circ C$. Again, in the presence of base, only the 3,5-dibromo betaine **5** was obtained (75%).

Other electrophiles were tested against **1**. Thus, phenyldiazonium chloride produced the diazo compound **6** as a red solid (50%). Reactions with Michael substrates were not successful, due to the good leaving




Fig. 2 Reagents and Conditions: i) MeI/acetone, r.t. or BzCl/toluene, reflux; ii) $\text{I}_2/\text{CH}_2\text{Cl}_2$; iii) $3\text{I}_2/\text{K}_2\text{CO}_3/\text{CH}_2\text{Cl}_2$, r.t.; iv) $2\text{Br}_2/\text{K}_2\text{CO}_3/\text{CH}_2\text{Cl}_2$, r.t.; v) $\text{PhN}_2^+ \text{Cl}^-/\text{water}$, r.t.; vi) Quinone/silica/ CH_2Cl_2 , r.t.

character of the aminide, but with quinones the initial adduct eliminated pyridine as described in a related process by Kakehi,⁵ when silica gel was used as catalyst. Thus quinone adducts **7a** (64%) and **7b** (65%) were obtained. Finally, cleavage of the N-N bond was carried out using a Zn/acetic acid reduction system,⁶ and in this way the corresponding 2-aminopyridines **8** were obtained with yields.

In summary, the use of heteroaryl-stabilized azinium aminides provides an easy, mild and selective approach to differently substituted 2-aminopyridines. Further experiments are in progress to extend the process as a general methodology to other azines.

Acknowledgements. Authors wish to thank the Comisión Interministerial de Ciencia y Tecnología (C.I.C.Y.T.) for financial support (Proj. PB 90-0284), Ministerio de Educación y Ciencia (M.E.C.) for a grant to one of us (R.C.M.) and ALTER S.A. for a generous gift of chemicals.

References.

1. a) Ollis, W. D.; Stanford, S. P.; Ramsden, C. A. *Tetrahedron* **1985**, *41*, 2239. b) Tamura, Y.; Ikeda, M. *Adv. Heterocyclic Chem.* **1981**, *29*, 71. c) Tsuge, O., Kanemasa, S. *Adv. Heterocyclic Chem.* **1989**, *45*, 232. d) Katritzky, A. R.; Lam, J. N. *Heterocycles* **1992**, *33*, 1011 on N-Oxides and N-Imides.
2. Cuadro, A. M., Novella, J. L., Molina, A., Alvarez-Builla, J., Vaquero, J. J. *Tetrahedron* **1990**, *46*, 6033.
3. Beyer, H., Thieme, E. *J. Prakt. Chem.* **1966**, *31*, 293.
4. Spitsin, N. V., Kochkanyan, R. O. *Kim. Geterotsikl. Soedin.* **1989**, 626.
5. Sasaki, T., Kanematsu, K., Kakehi, A. *Tetrahedron* **1972**, *28*, 1469.
6. Timpe, H.-J., *Z. Chem.* **1972**, *12*, 333.

(Received in UK 22 January 1993)