

Document downloaded from the institutional repository of the University of Alcalá: <http://dspace.uah.es/dspace/>

This is a postprint version of the following published document:

Fonseca, W., Rey Benayas, J.M. & Alice, F.E., 2011. Carbon accumulation in the biomass and soil of different aged secondary forests in the humid tropics of Costa Rica. *Forest Ecology and Management*, 262(8), pp.1400–1408..

Available at <https://doi.org/10.1016/j.foreco.2011.06.036>

© 2011 Elsevier

Universidad
de Alcalá

(Article begins on next page)



This work is licensed under a
Creative Commons Attribution-NonCommercial-NoDerivatives
4.0 International License.

1

2 **Carbon accumulation in the biomass and soil of different aged secondary forests in the Humid Tropics**
3 **of Costa Rica**

4

5 William Fonseca¹, José María Rey Benayas², Federico E. Alice³

6

7 ¹Corresponding author. Escuela de Ciencias Ambientales, Universidad Nacional de Costa Rica, Campus

8 Omar Dengo 86-3000, Heredia, Costa Rica. wfonseca@una.ac.cr

9 ²Departamento de Ecología, Universidad de Alcalá, España. e-mail: josem.rey@uah.es

10 ³Escuela de Ciencias Ambientales, Universidad Nacional de Costa Rica, Heredia, Costa Rica.

11 falice@costarricense.cr

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

1 **Abstract**

2 Efforts are needed in order to increase confidence for carbon accounts in the land use sector, especially in
3 tropical forest ecosystems that often need to turn to default values given the lack of precise and reliable site
4 specific data to quantify their carbon sequestration and storage capacity. The aim of this study was then to
5 estimate biomass and carbon accumulation in young secondary forests, from 4 and up to 20 years of age, as
6 well as its distribution among the different pools (tree including roots, herbaceous understory, dead wood,
7 litter and soil), in humid tropical forests of Costa Rica. Carbon fraction for the different pools and tree
8 components (stem, branches, leaves and roots) was estimated and varies between 37.3 (\pm 3.3) and 50.3% (\pm
9 2.9). Average carbon content in the soil was 4.1% (\pm 2.1). Average forest plant biomass was 82.2 (\pm 47.9) Mg
10 ha⁻¹ and the mean annual increment for carbon in the biomass was 4.2 Mg ha⁻¹ year⁻¹. Approximately 65.2%
11 of total biomass was found in the aboveground tree components, while 14.2% was found in structural roots
12 and the rest in the herbaceous vegetation and necromass. Carbon in the soil increased by 1.1 Mg ha⁻¹ year⁻¹.
13 Total stored carbon in the forest was 180.4 Mg ha⁻¹ at the age of 20 years. In these forests, most of the carbon
14 (51-83%) was stored in the soil. Models selected to estimate biomass and carbon in trees as predicted by basal
15 area had R² adjustments above 95%. Results from this study were then compared with those obtained for a
16 variety of secondary and primary forests in different Latin-American tropical ecosystems and in tree
17 plantations in the same study area.

18

19 *Key words:* biomass models, carbon pools, tree plantations, natural regeneration, succession age.

20

21 **1. Introduction**

22

23 Growing forests and tree plantations and their soils are major sinks of atmospheric carbon (FAO, 2006; IPCC,
24 2007; Saugier and Pontailier, 2006; Schimel et al., 2001), and thus the influence of forests in the global
25 carbon cycle is now widely recognized (Basu, 2009; Bonan, 2008; González et al., 2008). Forest vegetation
26 captures atmospheric CO₂ through photosynthesis and stores it mainly in hard biomass (wood) with a slow
27 turnover rate of 14 -19 years for native forests in Chile (Gayoso and Guerra, 2005), around 50 to 100 years in
28 the Amazon (Vieira et al., 2005) and an average of 50 years according to Reeburgh (1997). This rate has been

1 estimated for one to two decades for secondary forests in Puerto Rico when considering litter (Ostertag et al.,
2 2008). Atmospheric carbon incorporation rates into the biomass or soil tend to decrease with forest age, being
3 it higher at young or intermediate ages (Gayoso and Guerra, 2005; Ostertag et al., 2008; Saynes et al., 2005).
4 Forests also mobilize atmospheric carbon through plant respiration and organic material decomposition,
5 although these losses are usually less than the gains. An exception is old growth forests or forests suffering
6 from acute degradation, where losses can exceed gains (CATIE, 2002). Forests, in addition, may transfer
7 organic material towards the water table or groundwater or other aquatic ecosystems (FAO, 2002; Percy et al.,
8 2003).

9
10 The world's forest cover is now around 4 billion ha (0.59 ha per capita) (FAO, 2009). Secondary forests
11 (those regenerating largely through natural processes after significant human and/or natural disturbance of the
12 original forest vegetation; Chokkalingam and de Jong, 2001) represent 35% of the tropical forests (Emrich et
13 al., 2000), approximately 850 million hectares (FAO, 2006), but accounts on land area under this type of
14 forest cover are hard to assess. For example, in Costa Rica, the area of secondary forests under different
15 succession stages is uncertain and several estimates have been provided during the last years. Joyce (2006)
16 provided an estimate of 793 811 ha in 2004, according to MINAE-SINAC (2007) there are 586 967 ha and in
17 the most recent study, up to 900 000 ha were found (Costa Rica, 2010). In any case, it is likely that this
18 ecosystem is increasing its cover promoted by the instability of prices from agricultural products and the
19 migration of inhabitants from rural areas to more urban areas (Aide and Grau, 2004; Grau and Aide, 2008;
20 Rey Benayas, 2005).

21
22 Adding to this uncertainty, when accounting for the carbon absorption and storage capacity of forest
23 ecosystems, many authors agree on the weaknesses from current estimates (Chave et al., 2004; Sarmiento et
24 al., 2005). Given the lack of site specific data, these estimates have to be performed using generic values on
25 the amounts of biomass, carbon in the biomass or generic allometric equations to determine biomass and
26 carbon for a given forest ecosystem. This procedure will hardly reflect in these accounts the interactions
27 between environmental and anthropogenic factors that cause variations in the carbon concentrations within
28 the biomass (with global variations ranging from 1 to 35 t CO₂ ha⁻¹ yr⁻¹; IPCC, 2007) (Sarmiento et al.,

1 2005; Keith et al., 2009) and a range of estimated emissions from the land-use change as wide as 0.5–2.7 GtC
2 for the 1990s (Ravindranath et al., 2007).

3
4 In the Latin American context, the majority of studies document growth in biomass and carbon storage in
5 primary forests, mainly in the woody material (Acosta et al., 2002; Schlegel et al., 2001; Segura et al., 2000).
6 Studies on secondary forests are scarcer (Feldpausch et al., 2007; Fonseca et al., 2008; Herrera et al., 2001)
7 and even more, the quantification of total plant biomass and carbon, including roots, has not been a common
8 practice, and is even rarer for secondary forests.

9
10 Under this context, we conducted a research in the Costa Rican Caribbean region, with the aim of estimating
11 the amount of biomass and carbon accumulated and stored in young secondary forests, as well as its
12 distribution among the different pools (tree, herbaceous vegetation, necromass, and soil). Since precise
13 estimations of all biomass and carbon pools are expensive and time consuming, we developed models to
14 estimate biomass and carbon stored by area unit, so simple field measurements allow for these estimations at
15 these ecosystems in the future. In addition, the carbon fraction in the biomass was determined for the different
16 components of the biomass.

17 18 **2. Materials and Methods**

19 **2.1 Study Area**

20 This research was developed in the Costa Rican Caribbean region, which corresponds to a very humid tropical
21 forest life zone, according to Holdridge's Life Zone classification system (1967). The altitude ranges between
22 50 and 350m asl. Predominating climate is humid to very humid, hot to very hot, with or without a dry season
23 of < 25 intermittent days with water deficit per year (Herrera, 1985; Mena, 2007). The mean annual
24 precipitation varies between 3420 and 6840 mm and mean annual temperature between 25 and 27°C. Forests
25 are found on soils that are Ultisols and Inceptisols, with < 35% base saturation, these are deep, well drained,
26 red or yellow in color and with relatively low fertility. Both of these soil types are located on land with slopes
27 that range between 2% and 15% (ITCR, 2004).

1 **2.2 Establishment of sampling plots**

2 Seven sites were selected within the study area with secondary forests that range between 4 to 20 years of age.
3 Selection criteria was based on access to the forests (landowners willingness to support research), landowners
4 knowledge of the forest age and an appropriate distribution and representativeness of ages. These forests
5 were therefore found in private lands which were mostly abandoned pasture lands and for which age was
6 determined based on the landowner's knowledge of land abandonment. In each site, two to six 500-m²
7 rectangular sampling plots were established to estimate forest biomass. The number of plots at each site
8 depended on the variation of the secondary forest regeneration age (at least one plot per identified age) and
9 the heterogeneity of the secondary forest (i.e., if one coetaneous secondary forest showed a heterogeneous
10 vegetation structure, this was measured through the establishment of two or more plots). A total of 38 plots
11 were established, out of which 10 plots were re-measured two years after the first measurement in order to
12 complete an appropriate age distribution for a total of 48 plots sampled. Some forests with similar ages were
13 grouped to simplify analysis. These correspond to ages 4.5 and 6.5 (Table 2), which compile data averaged
14 for ages 4 - 5 and 6 - 7 accordingly.

15

16 Each 500-m² plots included four 1-m² and one 5-m² subplots to sample particular biomass compartments (see
17 below). In addition, 11 plots that represented a baseline from which secondary succession started were also
18 sampled. These sites were within the farms, adjacent to secondary forests sampled and where the current land
19 use is still pasture land. Baseline vegetation consisted mostly of grasses from the Poaceae family.

20

21 **2.3 Biomass Estimation**

22 *2.3.1 Aboveground tree biomass*

23 Aboveground tree biomass is usually determined through the selection of a single tree based on the dbh.
24 MacDicken (1997) recommends the selection of a tree with mean basal area, Schlegel et al. (2001)
25 recommends the random selection of one tree per diametric class of the most abundant species in each class.
26 For this study, the selection of trees to be harvested was based on the Importance Value Index (IVI), the IVI
27 being the sum of abundance, frequency and dominance or basal area expressed in relative values (Krebs,
28 1985). In each plot, every individual was classified into 5cm interval diametric classes, and the species with

1 the highest Importance Values Index (IVI) were determined for each class. In each sampling plot, all woody
2 plants with a diameter at breast height (dbh) ≥ 2.5 cm were measured. These accounted for a total of 6984
3 individuals, which were identified to the species (66.6%), genera (30.3%) and family level (0.38%) or
4 remained unknown (2.7%). In each plot, every individual was classified into 5cm interval diametric classes,
5 and selected a mean tree for each class as a sample. A total number of 193 trees corresponding to 35 different
6 species whose diameter varied between 2.8 and 28.2cm were sampled. The biomass was determined through
7 field measurements for weight for each tree component (leaves, branches and stem).

8 9 *2.3.2 Belowground tree biomass*

10 Belowground tree biomass in this study mainly refers to the structural or "anchor" roots and all of the fine
11 roots attached to the main root after harvesting. Roots with a diameter > 5 mm (according to the classification
12 proposed by Sierra et al. (2001)) were estimated through the excavation and extraction of the root system for
13 the average selected trees. Excavation and extraction was carried out with a retro-excavator or trencher,
14 agricultural tractor and/or manually with a chain hoist. These roots were then washed in the field and weighed
15 once they were air dry for one – two hrs.

16 17 *2.3.3 Biomass in herbaceous vegetation, small woody material and seedlings*

18 Grasses, lianas, ferns, shrubs and some tree seedlings with a dbh < 2.5 cm, were measured in 1 x 1m subplots
19 located in every corner of the main 500 m² plot. In each 1 m² subplot all plant material was harvested to
20 ground level, all four subplots were grouped into one sample and weighed in the field.

21 22 *2.3.4 Necromass*

23 Necromass or dead woody material found at ground level was divided into fine necromass (litter and woody
24 material < 2 cm in diameter) and large necromass (dead woody material ≥ 2 cm in diameter). Fine necromass
25 was estimated at four 0.5 x 0.5 m subplots that were randomly distributed throughout the 500m² plot; these
26 four samples were grouped into one sample for analysis. Large necromass was estimated at one 5 x 5 m
27 subplot that was randomly placed within the 500 m² plot. The collected material was weighed in the field.

1 **2.4 Soil organic carbon and carbon fraction in the biomass**

2 The total amount of carbon stored in the soil was quantified based on the soil's carbon content, bulk density
3 and sampling depth. A total of four 30-cm depth soil samples were randomly selected within each main plot,
4 extracted and mixed together in order to obtain a sample of approximately 1 kg. Bulk density was determined
5 through the cylinder method (MacDicken, 1997), collecting one cylinder per plot.

6
7 **2.5 Carbon fraction analysis in plant material and soil**

8 For every sample weighed in the field, an approximately 1 kg sub-sample was collected and taken to the
9 laboratory in order to determine the carbon fraction. Each sub-sample of the different components of the
10 biomass was taken to the lab and dried in an oven at 60°C for 72 hours to estimate its dry matter content
11 (DMC). Soil samples were dried at 55°C for three days and subsequently ground and run through a 240-µm
12 sieve. Carbon content in the plant biomass and soil was determined following the methods by Pregl and
13 Dumas (Bremner and Mulvaney, 1982) in an auto-analyzer (Perkin-Elmer series II, CHN/S 2400, Norway
14 Co.).

15
16 **2.6 Increases in the carbon content in plant biomass and soils**

17 The Mean Annual Increment (MAI) was calculated for the biomass and for the carbon in the biomass as MAI
18 = B or C/t, where B is biomass, C is carbon, both expressed in Mg ha⁻¹, and t is the number of years. In order
19 to analyze changes in the carbon content from soils, a current annual increment (CAI) was estimated for each
20 age, being it the difference between two periods and the number of years for that period (ICM= ΔC/t)
21 (Prodan et al.,1997). In this case, an average of the carbon content found in the soil at a certain age minus the
22 average carbon content in the baseline for that age class divided by the number of years.

23
24 **2.7 Models to estimate biomass and carbon**

25 Models were adjusted using the method of ordinary least squares (Fonseca et al., 2009). Approximately 25
26 models were tested for total tree biomass (Mg ha⁻¹), total forest biomass (Mg ha⁻¹) and total carbon in the
27 biomass (Mg ha⁻¹). The methodology presented by Salas (2002) and Segura and Andrade (2008) was followed
28 in order to determine the best fit equation. The selected models with logarithmic transformations were later

1 corrected using a correction factor (CF) as explained by Sprugel (1983). The suggested equation to estimate
2 the correction factor is: $CF = \exp(SSE^2/2)$, where: SSE = estimated standard error by the regression.

3 4 **3. Results**

5 **3.1 Carbon fraction in the biomass and soil carbon content**

6 The carbon fraction for the more lignified biomass components (stem, branches, roots and large necromass) in
7 secondary succession of 4 to 20 years, varied between 47.3 (± 3.3) and 50.3% (± 2.9). The carbon fraction for
8 leaves, herbaceous vegetation and fine necromass (litter) varied between 37.3 (± 3.3) and 43.5% (± 1.9). The
9 average across biomass compartments was 46.8% ± 4.0 . Soil had a carbon content of 4.1% (± 1.9 ; Table 1).

10 11 **3.2 Carbon accumulation in plant biomass**

12 Total biomass increased with the age of the secondary succession and had a positive correlation (Figure 1).
13 The increase in biomass was fast during the first 10 years, when total biomass averaged 174.5 \pm 16.4 Mg ha⁻¹.
14 It then decreased with lower values found in secondary forests older than that age (Figure 1; Table 2). The
15 average total biomass for all different ages was 82.2 \pm 47.9 Mg ha⁻¹. The mean annual increments for total
16 biomass and carbon in the biomass were 8.9 and 5.3 Mg ha⁻¹ year⁻¹, respectively.

17
18 Most of the carbon in the forest plant biomass was stored in the trees, with an average of 80.1 \pm 15.3% at all
19 different ages, followed by necromass (15.8 \pm 13.0%) and herbaceous vegetation (4.2 \pm 5.5%) (Figure 1; Table
20 2). There were no significant differences between the amounts of biomass and carbon in the biomass in
21 compartments other than trees (Table 2). Within the tree compartments, biomass and carbon were highest in
22 the stem (58.4 \pm 11.8%), and lower in the branches (18 to belowground ratio was marginal and negatively
23 correlated with forest age ($r = -0.27$, $P = 0.06$, $n = 48$).

24 25 **3.3 Carbon accumulation in the soil**

26 The amount of carbon in the soil was positively correlated with the age of the secondary forest, but this was a
27 relatively weak relationship (Figure 1). The increase of carbon in the soil was 1.09 Mg ha⁻¹ year⁻¹. At all ages,

1 the amount of carbon accumulated in the soil was higher than the amount of carbon stored in total biomass
2 (Figure 1; Table 2).

3
4 At the forest level (biomass and soil), the carbon accumulated increased with forest age (Figure 1; Table 2). In
5 recently established forests, where the biomass mainly corresponded to herbaceous vegetation, 98.8% of the
6 total carbon was stored in the soil. However, the relative amount of the total forest carbon in the soil
7 decreased rapidly as the succession progressed on, and 73.77% of total carbon was stored in the soil in 20-
8 year old forests (Figure 1).

9

10 **3.4 Biomass and Carbon Models**

11 Three models were selected to estimate biomass and carbon in the biomass that were highly predictive (Table
12 4). These models had R^2 adjustments above 95% and the significance levels was $p < 0.01$; the models' standard
13 errors were low and showed a normal distribution. The use of a correction factor (CF) increases the amount of
14 estimated biomass and carbon by $< 5\%$. The models that used age and diameter at breast height as predictive
15 variables of total biomass and carbon did not show good adjustments (results not shown).

16

17 **4. Discussion**

18

19 **4.1 Carbon content in the biomass of secondary forests**

20 In this study we estimated the amount of accumulated carbon in the biomass for the different components of
21 young secondary forests of the Costa Rica Caribbean Region. Few previous studies have estimated the precise
22 values of the amount of carbon found in the biomass of secondary forest species, and thus to transform the
23 amount of dry biomass into carbon a 0.5 conversion factor (Hoen and Solberg, 1994; Husch, 2001; Sarmiento
24 et al., 2005) is generally used. For the studied forests, the lowest carbon fractions of plant biomass
25 corresponded to those components with less lignin, such as fine necromass, leaves and herbaceous vegetation,
26 while large necromass, stem, roots and branches had higher carbon concentrations (Gayoso and Guerra, 2005;
27 Gifford, 2000). Other studies have not found differences in the carbon content of the different tree
28 components (Segura et al., 2000). Furthermore, higher carbon concentrations in components such as leaves

1 have been reported (Gifford, 2000). The results obtained in this study are found within the limits reported for
2 forest plantations in the same region (Cubero and Rojas, 1999; Fonseca et al., 2009).

4 **4.2 Carbon accumulation in secondary forests**

5 The maximum biomass accumulation in this study and the mean annual increment (174.5 Mg ha^{-1} and 8.9 Mg
6 $\text{ha}^{-1} \text{ yr}^{-1}$, respectively) are found within the range reported for secondary tropical forests by other studies
7 (Chacón et al., 2007; Marín et al., 2007; Yan et al., 2007). The average MAI for biomass, excluding roots,
8 from all these studies was $7.83 \text{ Mg ha}^{-1} \text{ yr}^{-1}$.

9
10 We found that the amount of carbon stored in the soil represented 74.3% of the total carbon in the forest,
11 51.5% higher than the biomass. Other studies in tropical areas have found that the amount of soil carbon was
12 between 50 and 75% of the total forest carbon (Fonseca et al., 2008; Jandl, 2006; Lagos and Vanegas, 2003).

13
14 The average increase and the percentage of organic carbon in the soil reported by other studies, for primary
15 and secondary tropical forests, are $0.5\text{-}2.0 \text{ MgC ha}^{-1} \text{ yr}^{-1}$ and 3.34%, respectively (Feldpausch et al., 2007;
16 Fonseca et al., 2008; Liu et al., 2006), are similar to the values of $1.09 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ and 4.2% found in this
17 study. In spite of the contribution that soil plays in the ecosystem's total carbon, approximately 75% of the
18 references cited by this study did not evaluate this carbon pool, due to its difficulty and high costs (Brown et
19 al., 1989; de Jong et al., 2000; MacDicken, 1997).

20
21 There is some controversy with regards to the role of land use change and the accumulation of carbon in the
22 soil, where previous land use is said to be a determining factor in either soil carbon accumulation or loss (Post
23 and Kwon 2000, Guo and Gifford, 2002; IPCC, 2007). Considering that pasture lands contain a higher
24 amount of fine roots that incorporate more carbon to the soil due to a fast decomposition rate as opposed to a
25 more lignified radical system from trees, Guo and Gifford (2002) have reported for carbon loss from this pool
26 when changing from pastures to secondary forests. Other studies have found an increase in carbon and thus
27 recognize that soil carbon tends to increase as the succession moves forward due to the contribution of
28 organic matter from roots and decomposing detritus (Hughes et al., 1999; Powers and Veldkamp, 2005;

1 Schedlbauer and Kavanagh, 2008; Sierra et al., 2001; Veldkamp et al., 2003). Still, other studies have failed
2 to find differences among different age groups (Gamboa et al., 2008; Ostertag et al., 2008; Tschakert et al.,
3 2007).

4
5 Accordingly, we did find a positive but low correlation between the amount of soil carbon and the age of the
6 forest, in contrast with the high correlation found between biomass and forest age (Figure 1). The low
7 correlation between soil carbon and forest age can be attributed partly to the slow incorporation of carbon into
8 the soil (Gamboa et al., 2008; McGrath et al., 2001; Robert, 2002; Saynes et al., 2005; Singh et al., 2007;
9 Turner et al., 2005) together with the young age of the studied forests. However, as reported by other authors,
10 previous land use, the number of years under the previous land use, the stage of the succession, distance from
11 seed sources and intervention or management, among others (Hughes et al., 1999; Mesquita, 2000) may all be
12 factors, that individually or in a combination, determine the amount of carbon found at the soil. In this
13 particular case, given that these secondary forests have grown in degraded, over pastured and more compacted
14 lands, we believe that besides from the reasons reported by other studies, there might also be an effect by
15 differences on bulk density from these soils and those found under secondary forests. However, this
16 assumption still needs to be proven.

18 **4.3 Comparison with tree plantations**

19 Fonseca et al. (unpublished results) determined the carbon accumulated in the biomass and soil of managed
20 forest plantations of comparable ages (from 0 to 16 years of age), found within the same region and therefore
21 under similar conditions. Furthermore, these also followed the same land use pattern, changing from pasture
22 lands to forest lands. MAI values for these ecosystems were of 7.1 MgC ha⁻¹ yr⁻¹ in the biomass of *Vochysia*
23 *guatemalensis* and 5.3 MgC ha⁻¹ yr⁻¹ in *Hieronyma alchorneoides* plantations. For these same plantations,
24 increases from carbon in the soil were 1.7 and 1.3 Mg ha⁻¹ yr⁻¹ respectively. These results show how forest
25 plantations (without taking into account biomass from thinnings and in spite of these being four years
26 younger) have almost twice the ability to store carbon in the biomass when compared to secondary forests
27 under similar conditions and an approximately 20% more with regards to soil carbon. At an ecosystem level,
28 MAI for total carbon in forest plantations was of 8.7 and 6.5 MgC ha⁻¹ yr⁻¹, higher than the 5.3 MgC ha⁻¹ yr⁻¹

1 found in secondary forests from this study. This can be partly explained by the effect of silvicultural activities
2 aimed at increasing the amount and quality of forest productivity (Daniel et al., 1982; Kerr and Morgan, 2006;
3 Wadsworth, 2000), therefore increasing carbon accumulation and organic material incorporated to the soils.

4 5 **4.4 Models to estimate biomass and carbon**

6
7 In the tropics, models to estimate biomass and carbon tend to show good R^2 adjustments when relating the
8 biomass to diameter, basal area and/or height of individual trees (Fonseca et al., 2009). Most previously
9 published models have been used to estimate biomass and carbon per tree and/or tree component (Acosta et
10 al., 2002; Brandeis et al., 2006; Gaillard et al., 2002; Litton and Kauffman, 2008; Lagos and Vanegas, 2003;
11 Segura and Kanninen, 2005) but we have not found models to estimate biomass and carbon per area unit.
12 Where in doing so, our results from the tested models show a good adjustment ($R^2_{aj} > 95\%$) equal to or above
13 reported by other works, which we mainly attribute to our high number of samples distributed over a wide
14 range of ages. In this work, the tested models using age as the predicting variable for biomass and carbon did
15 not show good R^2 adjustments, a behavior that has been a common trend in other studies (Feldpausch et al.,
16 2004). Overall, the good adjustment of the selected models to estimate biomass and carbon per hectare using
17 simple field variables such as basal area represents an important advance towards the precise and reliable
18 quantification of carbon accumulation in tropical secondary forests.

19 20 **5. Acknowledgements**

21 The authors would like to express their sincere gratitude to Johan Montero and Henry Toruño, researchers at
22 the Forestry Research and Services Institute from the National University of Costa Rica, for their support
23 during field data collection. This work received finance from the National University of Costa Rica, the Costa
24 Rican Ministry of Science and Technology and from the private sector.

25 26 **6. References**

27 Acosta, M., Hernández, J., Velásquez, A., Etchevers, J., 2002. Estimación de la biomasa aérea mediante el
28 uso de relaciones alométricas en seis especies arbóreas en Oaxaca, México. *Agrocienca* 36(6):725-736.

1 Aide, T., Grau, H., 2004. Globalization, migration, and Latin American ecosystems. *Science* 305:1915-1916.

2 Basu, P., 2009. A green investment. If growing forests in India can generate lucrative carbon credits, then
3 why isn't everyone planting trees? News Feature. *Nature* 457(8):144-146.

4 Bonan, G.B., 2008. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests.
5 *Science* 320:1444-1449.

6 Brandeis, T., Delaney, M., Parresol, B., Royer, L., 2006. Development of equations for predicting Puerto
7 Rican subtropical dry forest biomass and volumen. *Forest Ecology and Management* 233: 133-142.

8 Bremner J.M., Mulvaney C., 1982. Carbon, inorganic nitrogen. In R. Miller and D. Keeney (eds.). *Methods*
9 *for soil analysis: chemical and microbiological properties*, pp 552, 673-682. 2 ed. American Society of
10 Agronomy, Madison, US.

11 Brown, S., Gillespie A., Lugo, A., 1989. Biomass estimation methods for tropical forests with applications to
12 forest inventory data. *Forest Science* 35(4):881-902.

13 Canadell, J., Kirschbaum, M., Kurz, W., Sanz, M.J., Schlamadinger, B., Yamagata, Y., 2007. Factoring out
14 natural and indirect human effects on terrestrial carbon sources and sinks. *Environ. Sci. Policy*
15 doi:10.1016/j.envsci.2007.01.009.

16 CATIE (Centro Agronómico de Investigación y Enseñanza, CR), 2002. Planificación del manejo
17 diversificado de bosques latifoliados húmedos tropicales. Lorena Orozco (Ed). Turrialba, Costa Rica,
18 CATIE. 329 p. (Serie Técnica/Manual Técnico n° 56).

19 Chacón, P., Leblanc, H., Russo, R., 2007. Fijación de carbono en bosques secundarios de la región tropical
20 húmeda en Costa Rica. *Tierra Tropical* 3(1):1-11.

21 Chave, J., Condit, R., Aguilar, S., Hernandez, A., Lao, S., Perez, R. 2004. Error propagation and scaling for
22 tropical forest biomass estimates. *Phil. Trans. R. Soc. Lond. B* 359: 409–420. [http://si-](http://si-pddr.si.edu/dspace/bitstream/10088/6729/1/Chave_Condit_Aguilar_Hernandez_Lao_and_Perez_2004.pdf)
23 [pddr.si.edu/dspace/bitstream/10088/6729/1/Chave_Condit_Aguilar_Hernandez_Lao_and_Perez_2004.p](http://si-pddr.si.edu/dspace/bitstream/10088/6729/1/Chave_Condit_Aguilar_Hernandez_Lao_and_Perez_2004.pdf)
24 [df](http://si-pddr.si.edu/dspace/bitstream/10088/6729/1/Chave_Condit_Aguilar_Hernandez_Lao_and_Perez_2004.pdf)

25 Chokkalingam U. and de Jong W. 2001. Secondary forest: a working definition and typology. *International*
26 *Forestry Review* 3(1): 19-26.

27 Costa Rica, 2010. Readiness Preparation Proposal, R-PP Costa Rica. Presented to the Forest Carbon
28 Partnership Facility (FCPF). FONAFIFO/MINAET.

1 <http://www.forestcarbonpartnership.org/fcp/sites/forestcarbonpartnership.org/files/Documents/PDF/Sep>
2 [2010/R-PP_Costa_Rica_English_08-19-10.pdf](http://www.forestcarbonpartnership.org/fcp/sites/forestcarbonpartnership.org/files/Documents/PDF/Sep)

3 Cubero, J., Rojas, S., 1999. Fijación de carbono en plantaciones de *Gmelina arborea*, *Tectona grandis* y
4 *Bombacopsis quinata*. Tesis de Licenciatura. Heredia, Costa Rica, Universidad Nacional, Escuela de
5 Ciencias Ambientales. 94 p. Available at:
6 [http://cglobal.imn.ac.cr/Pdf/mitigacion/Estudio%20sobre%20Fijacion%20de%20Carbono%20en%20Pla](http://cglobal.imn.ac.cr/Pdf/mitigacion/Estudio%20sobre%20Fijacion%20de%20Carbono%20en%20Plantaciones.pdf)
7 [ntaciones.pdf](http://cglobal.imn.ac.cr/Pdf/mitigacion/Estudio%20sobre%20Fijacion%20de%20Carbono%20en%20Plantaciones.pdf)

8 Daniel, P.; Helms, U.; Baker, F., 1982. Principios de silvicultura. I ed. McGraw-Hill. México, DF. 492 p.
9 de Jong, B., Tipper, R., Montoya, G., 2000. An economic analysis of the potential for carbon sequestration by
10 forest: evidence from southern México. *Ecological Economics* 33:313-327.

11 Emrich, A., Pokorny, B., Sepp, C., 2000. Investigación de los bosques: Importancia del manejo de los
12 bosques secundarios para la política de desarrollo bosques tropicales. ECO – Society for socio-
13 ecological programme consultancy Anette Emrich. Eschborn. 98 p + annexes.

14 FAO (Organización de la Naciones Unidas para la Agricultura y la Alimentación, It), 2002. Captura de
15 carbono en los suelos para un mejor manejo de la tierra. Basado en el trabajo de Michael Robert. Institut
16 National the Recherche Agronomique. Paris, Francia. 62 p.

17 FAO (Organización de la Naciones Unidas para la Agricultura y la Alimentación, It), 2006. Global Forest
18 Resource Assessments 2005: Progress towards sustainable forest management. FAO Forestry Paper
19 n°146. Consulted May 22 2006. Available at:
20 http://www.fao.org/documents/show_cdr.asp?url_file=/docrep/008/a0400e/a0400e00.htm

21 FAO (Food and Agriculture Organization of the United Nations, IT), 2009. State of the World's Forests.
22 Rome, Italy. ISBN 978-92-5-106057-5. 152 p.

23 Feldpausch, T., Rondon, M., Fernandes, E., Riha, S., Wandelli, E., 2004. Carbon and nutrient accumulation in
24 secondary forest regenerating from degraded pastures in Central Amazonia, Brazil. *Ecological*
25 *Applications* 14:164-176.

26 Feldpausch, T., Prates, C., Fernandes, E., Riha, S., 2007. Secondary forest growth deviation from
27 chronosequence predictions in central Amazonia. *Global Change Biology* 13(5):967-979.

- 1 Fonseca, W., Alice, F., Montero, J., Toruño, H., Leblanc, H., 2008. Acumulación de biomasa y carbono en
2 bosques secundarios y en plantaciones forestales de *Vochysia guatemalensis* e *Hieronyma alchorneoides*
3 en el Caribe de Costa Rica. *Agroforestería en las Américas* 46:57-63.
- 4 Fonseca, W., Alice, F., Rey Benayas, J.M., 2009. Modelos para estimar la biomasa de especies nativas en
5 plantaciones y bosques secundarios en la zona Caribe de Costa Rica. *Bosque* 30(1):36-47.
- 6 Fonseca, W., Alice, F., Rey Benayas, J.M., (Unpublished results). Site and species specific biomass
7 accumulation and carbon sequestration data for forest plantations of *Vochysia guatemalensis* and
8 *Hieronyma alchorneoides*.
- 9 Gaillard, C., Pece, M., Juárez, M., 2002. Biomasa aérea de quebracho blanco (*Aspidosperma quebracho-*
10 *blanco*) en dos localidades del Parque Chaqueño Seco. *Revista Ciencias Ambientales - Quebracho*
11 9:116-127.
- 12 Gamboa, A., Hidalgo, C., de León, F., Etchevers, J., Gallardo, J., Campo, J., 2008. Nutrient addition
13 differentially affects soil carbon sequestration in secondary tropical dry forests: Early- versus late-
14 succession stages. *Restoration Ecology* 18(2):252-260.
- 15 Gayoso, J., Guerra, J., 2005. Contenido de carbono en la biomasa aérea de bosques nativos en Chile. *Bosque*
16 26(2):33-38.
- 17 Gifford, R., 2000. Carbon contents of above-ground tissues of forest and woodland trees. Canberra:
18 Australian Greenhouse Office, National Carbon Accounting System, 2000, Technical Report n° 22. 17 p.
- 19 González, P., Hassan, R., Lakada, P., McCallum, I., Nilson, S., Pulhin, J., van Rosenburg, B., Scholes, B.,
20 2008. Forest and woodland systems: Chapter 21. Pp 587-614. Consulted Feb. 12 2009. Available at:
21 <http://www.millenniumassessment.org/documents/document.290.aspx.pdf>
- 22 Grau, R., Aide, M., 2008. Globalization and Land-Use Transitions in Latin America. *Ecology and Society*
23 13(2): art. 16. Available at: <http://www.ecologyandsociety.org/vol13/iss2/art16/>
- 24 Guo, L.B and Gifford L.M., 2002. Soil carbon stocks and land use change: A meta Analysis. *Global Change*
25 *Biology* 8(4): 345-360.
- 26 Herrera, M.A., del Valle, J.I., Orrego, S., 2001. Biomasa de la vegetación arbórea y leñosa pequeña y
27 necromasa en bosques tropicales primarios y secundarios de Colombia. Valdivia, Chile. Simposio

1 Internacional Medición y Monitoreo de la Captura de Carbono en Ecosistemas Forestales, 18 al 20 de
2 Octubre del 2001. 19 p.

3 Herrera, W., 1985. Clima de Costa Rica: Vegetación y Clima de Costa Rica. Volumen 2. Editor Luis Diego
4 Gómez. UNED, San José, Costa Rica. 118 p.

5 Hoen H., Solberg B., 1994. Potential and economic efficiency of carbon sequestration in forest biomass
6 through silvicultural management. *Forest Sci.* 40: 429-451.

7 Holdridge, L., 1967. *Life Zone Ecology*. San José, Costa Rica. Centro Científico Tropical. 82 p.

8 Hughes, R.F., Kauffman, J.B., Jaramillo, V.J., 1999. Biomass, carbon, and nutrient dynamics of secondary
9 forests in a humid region of México. *Ecology* 80(6):1892-1907.

10 Husch, B., 2001. Estimación del contenido de carbono en los bosques. Valdivia, Chile. Simposio
11 Internacional Medición y Monitoreo de la Captura de Carbono en Ecosistemas Forestales, 18 al 20 de
12 Octubre del 2001. 9 p. Available at: <http://www.uach.cl/simposiocarbono/doc/Husch.PDF>

13 IPCC (Intergovernmental Panel on Climate Change), 2007. Informe de síntesis. Contribución de los Grupos
14 de trabajo I, II y III al Cuarto Informe de evaluación del Grupo Intergubernamental de Expertos sobre el
15 Cambio Climático. Ginebra, Suiza, IPCC. 104 p.

16 ITCR (Instituto Tecnológico de Costa Rica), 2004. Atlas digital de Costa Rica. Laboratorio de Sistemas de
17 Información Geográfica, Escuela de Ingeniería Forestal. Cartago, CR.

18 Jandl, R., 2006. Secuestro de carbono en bosques: el papel del suelo. Taller internacional sobre secuestro de
19 carbono. IUFRO-RIFALC. 9 p.

20 Joyce, A.T., 2006. Land use change in Costa Rica: 1966 - 2006, as influenced by social, economic, political,
21 and environmental factors. San José, Costa Rica. Litografía e Imprenta LIL, S.A. 276 p. ISBN: 9977-47-
22 360-9.

23 Keith, H., Mackey, B.G., Lindenmayer, D.B. 2009. Re-evaluation of forest biomass carbon stocks and lessons
24 from the world's most carbon-dense forests. *PNAS*. 106 (28): 11635-11640.
25 <http://www.pnas.org/content/106/28/11635.short>.

26 Kerr, G.; Morgan, G., 2006. Does formative pruning improve the form of broadleaved trees? *Canadian*
27 *Journal of Forest Research* 36(1):132.141.

1 Krebs, J.C., 1985. Ecología: Estudio de distribución y abundancia. Segunda edición. Ed. Harla México. 753
2 p.

3 Lagos, O., Vanegas, S., 2003. Impacto del aprovechamiento forestal en la biomasa y carbono de bosques
4 naturales de Nueva Quezada, Río San Juan. Tesis Mag. Sc. Managua, Nicaragua, Universidad
5 Centroamericana. 121 p. Available at: [http://orton.catie.ac.cr/cgi-](http://orton.catie.ac.cr/cgi-bin/wxis.exe/?IsisScript=carbo.xis&expresion=Bosques&formato=1&cantidad=50)
6 [bin/wxis.exe/?IsisScript=carbo.xis&expresion=Bosques&formato=1&cantidad=50](http://orton.catie.ac.cr/cgi-bin/wxis.exe/?IsisScript=carbo.xis&expresion=Bosques&formato=1&cantidad=50)

7 Litton, C., Kauffman, B., 2008. Allometric Models for Predicting Aboveground Biomass in Two Widespread
8 Woody Plants in Hawaii. *Biotropica* 40(3): 313-320 2008.

9 Liu, J.X., Liu, S.G., Loveland, T.R., 2006. Temporal evolution of carbon budgets of the Appalachian forests
10 in the US from 1972 to 2000. *Forest Ecology and Management* 222(1-3):191-201.

11 MacDicken, K., 1997. A guide to monitoring carbon storage in forestry and agroforestry projects. Forest
12 carbon Monitoring Program. Winrock International Institute for Agricultural Development (WRI). 87 p.
13 Consulted Aug. 25 2007. Available at: <http://www.winrock.org/REEP/PUBSS.html>.

14 Marín, E., Silver, W.L., Ostertag, R., 2007. Long-term patterns in tropical reforestation: Plant community
15 composition and aboveground biomass accumulation. *Ecological Applications* 17(3):828-839.

16 McGrath, D., Smith, C., Gholz, H., Oliveira, F., 2001. Effects of land-use change on soil nutrient dynamics in
17 Amazonia. *Ecosystems* 4: 625-645.

18 Mena, M., 2007. Clima de Costa Rica, Vertiente del Caribe. Instituto Meteorológico Nacional. Consulted Jan.
19 3 2007. Available at: http://www.imn.ac.cr/educacion/climacr/vertient_caribe.html.

20 Mesquita, R.C.G., 2000. Management of advanced regeneration in secondary forests of the Brazilian
21 Amazon. *Forest Ecology and Management* 130:131-140.

22 MINAE-SINAC, 2007. Estrategia para la sostenibilidad de la producción forestal en Costa Rica periodo 2006
23 – 2010. 35 p.

24 Ostertag, R., Marín, E., Silver, W., Schulten, J., 2008. Litterfall and decomposition in relation to soil carbon
25 pools along a secondary forest chronosequence in Puerto Rico. *Ecosystems* 11(5):701-714.

26 Percy, K.E., Jarndl, R., May, J.P., Lavigne, M., 2003. El papel de los bosques en ciclo, la captura y el
27 almacenamiento de carbono. Boletín nº 1: Bosques y el ciclo de carbono e nivel mundial: fuentes y
28 sumideros. 6 p. Consulted Mar. 12 2009. Available at: <http://iufro.boku.ac.at/iufro/taskforce/hptfcs.htm>.

- 1 Post, W., Kwon, K. 2000. Soil carbon sequestration and land-use change; processes and potential. *Global*
2 *Change Biology* 6:317-327.
- 3 Powers, JS., Veldkamp, E., 2005. Regional variation in soil carbon and $\delta^{13}C$ in forests and pastures of
4 northeastern Costa Rica. *Biogeochemistry* 72:315-336.
- 5 Prodan, M., Peters, R., Cox, F., Real, P., 1997. *Mensura forestal. Serie de investigación y evaluación en*
6 *desarrollo sostenible. San José, Costa Rica. IICA, GTZ. 561 p.*
- 7 Ravindranath N. H. and Ostwald M. 2007. *Carbon Inventory Methods Handbook for Greenhouse Gas*
8 *Inventory, Carbon Mitigation and Roundwood Production Projects . Advances in Global Change*
9 *Research Volume 29, 2007, DOI: 10.1007/978-1-4020-6547-7*
- 10 Reeburgh, W.S., 1997. Figures summarizing the global cycles of biogeochemically important elements.
11 *Bulletin of the Ecological Society of America* 78:260-267.
- 12 Rey Benayas, J.M., 2005. Restoration after land abandonment. *In* S. Mansourian, D. Vallauri y N. Dudley,
13 editores. *Forests restoration in landscapes: beyond planting trees. Springer-Verlag, Nueva York, EUA.*
14 *Pp 356-360.*
- 15 Robert, M., 2002. Captura de carbono en los suelos para un mejor manejo de la tierra. Informe sobre recursos
16 mundiales de suelos no. 96. FAO. Roma, Italia. 61 p.
- 17 Salas C., 2002. Ajuste y validación de ecuaciones de volumen para un relicto del bosque de Roble-Laurel-
18 *Lingue. Bosque* 23(2): 81-92.
- 19 Sarmiento G., Pinillos M., Garay I., 2005. Biomass Variability in Tropical American Lowland Rainforests.
20 *ECOTROPICOS* 18(1):1-20.
- 21 Saugier, B, Pontailier, J., 2006. El ciclo global del carbono y sus consecuencias en la fotosíntesis en el
22 *Altiplano boliviano. Ecología en Bolivia* 41(3):71-85.
- 23 Saynes, V., Hidalgo, C., Etchevers, J., Campo, J., 2005. Soil C and N dynamics in primary and secondary
24 *seasonally dry tropical forests in Mexico. Agriculture, Ecosystems and Environment . Applied Soil*
25 *Ecology* 29(3):282-289.
- 26 Schedlbauer J, Kavanagh K., 2008. Soil carbon dynamics in a chronosequence of secondary forests in
27 northeastern Costa Rica. *For. Ecol. Manage.* 255: 1326-1335.

- 1 Schimel, D.S., House, J.I., Hibbard, K.A., et al., 2001. Recent patterns and mechanisms of carbon exchange
2 by terrestrial ecosystems. *Nature* 414:169-172.
- 3 Schlegel, B., Gayoso, J., Guerra, J., 2001. Medición de la capacidad de captura de carbono en bosques de
4 Chile y promoción en el mercado mundial: Manual de procedimientos: Muestreos de biomasa forestal.
5 Valdivia, Chile, Universidad Austral de Chile. 24 p.
- 6 Segura M., Andrade H., 2008. ¿Cómo hacerlo? ¿Cómo construir modelos alométricos de volumen, biomasa o
7 carbono de especies leñosas perennes? *Agroforestería de las Américas* 46: 89-96.
- 8 Segura, M., Kanninen, M., Alfaro, M., Campos, J.J., 2000. Almacenamiento y fijación de carbono en bosques
9 de bajura de la zona atlántica de Costa Rica. *Revista Forestal Centroamericana* 30:23-28.
- 10 Segura, M., Kanninen, M., 2005. Allometric models for tree volume and total aboveground biomass in a
11 tropical humid forest in Costa Rica. *Biotrópica* 37(1):2-8.
- 12 Sierra, C., del Valle, J.I., Orrego, S., 2001. Ecuaciones de biomasa de raíces y sus tasas de acumulación en
13 bosques sucesionales y maduros tropicales en Colombia. Valdivia, Chile. Simposio Internacional
14 Medición y Monitoreo de la Captura de Carbono en Ecosistemas Forestales, 18 al 20 de octubre del
15 2001. 16 p. Available at: http://www.uach.cl/procarbono/pdf/simposio_carbono/18_Sierra.PDF
- 16 Singh, S.K., Singh, A.K., Sharma, B.K., Tarafdar J.C., 2007. Carbon stock and organic carbon dynamics in
17 soils of Rajasthan, India. *Journal of Arid Environments* 68 (3):408-421.
- 18 Sprugel, D.G., 1983. Correcting for bias in log-transformed allometric equations. *Ecology* 64(1): 209-210.
- 19 Tschakert, P., Coomes, O.T., Potvin, C., 2007. Indigenous livelihoods, slash-and-burn agriculture, and carbon
20 stocks in Eastern Panama. *Ecol. Econ.* 60: 807- 820.
- 21 Turner, J., Lambert, M.J., Jhonson, D.W., 2005. Experience with patterns of change in soil carbon resulting
22 from forest plantation establishment in eastern Australia. *Forest Ecology and Management* 220(1-
23 3):259-269.
- 24 Veldkamp, E., Becker, A., Schwendenmann, L., Clark, D., Schulte – Bisping, H., 2003. Substantial labile
25 carbon stocks and microbial activity in deeply weathered soils below a tropical wet forest. *Glob. Change*
26 *Biol.* 9:1171-1184.

- 1 Vieira, S., Trumbore, S., Camargo, P.B., Selhorst, D., Chambers, J.Q., Higuchi, N., Martinelli, L.A., 2005.
2 Slow growth rates of Amazonian trees: Consequences for carbon cycling. *Proceedings of the National*
3 *Academy of Sciences of the United States of America* 102(51):18502-18507.
- 4 Yan, J., Wang, Y., Zhou, G., Zhang, D., 2007. Estimates of soil respiration and net primary production of
5 three forests at different succession stages in South China. *Global Change Biology* 12(5):810-821.
- 6 Wadsworth, F.H., 2000. *Producción forestal para América Tropical. Manual de agricultura 710-s.* USDA,
7 Departamento de Agricultura de los Estados Unidos, Servicio Forestal. 603 p.
- 8




Figure 1. Carbon accumulation in secondary forests of different ages and its distribution in the biomass and soil.

Table 1. Carbon fraction (%) in the biomass and soil carbon content (%) in different compartments of young secondary forests in the Humid Tropics of Costa Rica.

Statistics	Stem	Branches	Leaves	Roots	Herbaceous Vegetation	Large Necromass	Fine Necromass	Soil
X	47.9	47.3	37.3	47.5	43.5	50.3	41.1	4.1
SD	3.9	3.3	3.3	4.4	1.9	2.9	2.4	2.1
n	193	191	193	193	58	48	48	59

X= average, SD = standard deviation, n = number of samples.

Table 2. Accumulated biomass and carbon (numbers in bold correspond to carbon) in secondary forests and their distribution in the different biomass compartments and soil. All expressed in $Mg\ ha^{-1} \pm$ standard deviation.

Age	Number of sampling plots	G (m2/ha)	Total biomass	Tree Biomass	Fine necromass	Herbaceous Vegetation	Large necromass	C total biomass	C soil	ICA soil
0	11		2.5±0.3	0.0±0.0	0.0±0.0	2.53±0.3	0.0±0.0			
				0.0±0.0	0.0±0.0	1.15±0.1	0.0±0.0	1.15±0.1	99.1±38.5	
4.5	5	8.6±5.3	44.9±38.1	37.3±36.6	3.6±2.0	2.2±1.4	1.9±1.8			
				13.2±11.2	1.9±0.8	1.1±0.4	0.9±5.1	17.1±12.0	95.1±12.5	1.6±0.7
6.5	12	9.8±5.0	40.5±16.7	32.8±16.9	2.1±1.5	2.8±2.6	2.9±4.5			
				15.7±7.8	0.8±0.6	1.2±1.1	1.7±2.2	19.4±8.0	96.0±21.6	0.6±0.5
8	4	15.1±8.1	86.9±36.3	79.1±39.4	4.1±0.6	1.9±2.9	1.9±1.4			
				33.0±17.7	1.7±0.2	0.8±1.2	1.0±0.8	36.4±16.5	90.1±11.1	0
10	6	21.4±8.5	174.5±16.4	127.5±47.7	6.7±3.0	1.2±0.8	39.0±45.6			
				65.4±26.6	3.0±1.5	0.5±0.3	18.2±20.4	87.1	90.4±11.2	0
12	15	18.7±4.8	104.0±37.0	88.2±37.3	6.3±3.6(b)	1.4±0.9	8.1±14.3			
				41.9±17.2	2.6±1.4	0.6±0.4	4.0±7.2	49.2±17.1	114.8±31.9	2.3±2.1
18	4	16.9±3.9	67.2±7.3	52.9±2.6	5.9±2.2	2.0±0.7	6.4±8.3			
				24.9±1.6	2.5±1.2	0.8±0.3	3.5±4.6	31.7±4.9	115.1±32.1	1.4±1.7

20	3	18.4±1.3	102.3±19.3	92.8±20.8	3.4±1.4	1.3±0.9	4.8±7.4				
				42.9±6.0	1.4±0.6	0.6±0.4	2.4±3.7	47.3±7.4	133.1±28.5	1.7±1.4	

Table 3. Biomass and carbon in the different components of total biomass, trees, and ecosystem.

Total Biomass Components								
Statistics	Total Biomass	Necromass	Herbaceous	Litter	Tree Biomass (%)			
	(Mg ha⁻¹)	(%)	Vegetation (%)	(%)	Leaves	Branches	Stem	Roots
n	48	48	48	48	48	48	48	48
X	81.98	8.97	4.51	7.10	4.77	14.35	46.13	14.17
SD	47.90	12.59	5.98	5.94	2.10	6.89	11.97	5.48
Carbon in tree biomass (%)								
Statistics					Leaves	Branches	Stem	Roots
n					48	48	48	48
X					5.32	18.37	58.38	17.93
SD					1.68	7.32	11.76	6.35
Carbon in the ecosystem								
Statistics	Ecosystem	Necromass	Herbaceous	Litter	Tree biomass (%)		Soil Carbon	
	(Mg ha⁻¹)	(%)	Vegetation (%)	(%)			(%)	
n	48	48	48	48	48		48	
X	143.16	2.87	0.68	1.35	20.82		74.29	
SD	36.94	5.33	0.67	0.83	10.48		11.98	

X= average, SD = standard deviation, n = number of samples. *= Probability 95 percent.

Table 4. Models selected for the estimation of biomass and carbon accumulated in secondary forests.

Model	R² aj (%)	SEE	N	CF
Bt= exp(1.06839 + 0.80802 *√G	95.7	0.310	48	1.05
CBa= exp(0.15004 + 0.800996 *√G	97.8	0.216	48	1.02
CBt= exp(0.272739 + 0.816253 *√G	96.0	0.299	48	1.05

Bt= Total biomass (Mg ha⁻¹), CBa= Carbon in the tree biomass (Mg ha⁻¹), CBt= Carbon in the total biomass (Mg ha⁻¹), G = basal area m²ha⁻¹, R²aj = adjusted coefficient of determination; exp (natural log base = 2.718271), n = sample size, SEE = model's standard error, CF= c

Carbon accumulation in the biomass and soil of different aged secondary forests in the Humid Tropics of Costa Rica

- We accounted for carbon accumulation in different aged secondary forests in the Costa Rican Caribbean region.
- All carbon pools were included in these estimates, where soils resulted in the most important carbon reservoir although not necessarily the most important in terms of carbon accumulation rates, this happened to be the aboveground biomass and most importantly the tree stem. However, other pools not regularly measured, happen to have an important contribution to the overall carbon accumulation rates and thus should be considered in future estimates.
- Biomass models to account for biomass and carbon in the different pools and at the ecosystem level, were created and its use recommended for sites under similar conditions.